

A Note on Semidensities in Antisymplectic Geometry

K. BERING¹

Institute for Theoretical Physics & Astrophysics
Masaryk University
Kotlářská 2
CZ-611 37 Brno
Czech Republic

February 1, 2008

Abstract

We revisit Khudaverdian's geometric construction of an odd nilpotent operator Δ_E that sends semidensities to semidensities on an antisymplectic manifold. We find a local formula for the Δ_E operator in arbitrary coordinates and we discuss its connection to Batalin-Vilkovisky quantization.

MCS number(s): 53A55; 58A50; 58C50; 81T70.

Keywords: Batalin-Vilkovisky Field-Antifield Formalism; Odd Laplacian; Antisymplectic Geometry; Semidensity; Halfdensity.

¹E-mail: bering@physics.muni.cz

1 Introduction

Recall that for a symplectic manifold with an even symplectic two-form $\omega = \frac{1}{2}dz^A\omega_{AB}dz^B$, there exists a canonical measure density given by the Pfaffian $\rho = \text{Pf}(\omega_{AB})$, *i.e.* there is a natural notion of volume in a symplectic manifold. A related fact is the Liouville Theorem, which states that Hamiltonian vector fields are divergenceless. On the other hand, the situation is completely different for an odd symplectic manifold, also known as an antisymplectic manifold and endowed with an odd antisymplectic two-form $E = \frac{1}{2}d\Gamma^A E_{AB}d\Gamma^B$. These geometries for instance show up in the Lagrangian quantization method of Batalin and Vilkovisky [1]. It turns out that there is *no* canonical choice of measure density ρ in this case, as, for instance, the above Pfaffian. This is tied to the fact that there is no meaningful notion of a superdeterminant/Berezinian for a matrix that is intrinsically Grassmann-odd. However, the upset runs deeper. In fact, a density ρ can never be a function of the antisymplectic matrix E_{AB} . Phrased differently, a density ρ always carries information that cannot be deduced from the antisymplectic structure E alone [2]. Within the standard Batalin-Vilkovisky framework, the possible choices of a density ρ is only partially determined by a requirement of gauge symmetry.

Around 1992 Batalin-Vilkovisky quantization took a more geometric form, in particular with the work of Schwarz [3]. The consensus was that the geometric setting requires two independent structures: an odd symplectic, non-degenerate two-form E and a measure density ρ . From these two structures, one may build a Grassmann-odd, second-order operator Δ_ρ , known as the odd Laplacian. Alternatively, one can view the odd Laplacian Δ_ρ itself as *the* fundamental structure of Batalin-Vilkovisky geometry [4, 5], which is conventionally required to be nilpotent.

Khudaverdian has constructed [6, 7, 8, 9] a Grassmann-odd, nilpotent, second-order operator Δ_E that does not rely on a choice of density ρ . The caveat is that the Δ_E operator is defined on semidensities rather than on scalars. (The notion of a semidensity is explained in eq. (3.1) below.) In retrospect, many pieces of Khudaverdian's construction were known to physicists, see for instance Ref. [10], p.440. In this short note we reconsider Khudaverdian's construction and find a local formula for the Δ_E operator that applies to arbitrary coordinate systems. The ability to work in any coordinates, not just Darboux coordinates, is important, since if one first has to search for a set of Darboux coordinates to the system that one is studying, symmetries (such as, *e.g.*, Lorentz covariance) or locality that one would like to preserve during the quantization process, are often lost.

The paper is organized as follows: We consider the antisymplectic structure in Section 2; the odd Laplacian Δ_ρ in Section 3; and in Sections 4 and 5, the Δ_E operator using Darboux coordinates and general coordinates, respectively. Finally, in Section 6 we analyze a modified Batalin-Vilkovisky scheme based on the Δ_E operator.

General remark about notation. We have two types of grading: A Grassmann grading ϵ and an exterior form degree p . The sign conventions are such that two exterior forms ξ and η , of Grassmann parity $\epsilon_\xi, \epsilon_\eta$ and exterior form degree p_ξ, p_η , respectively, commute in the following graded sense

$$\eta \wedge \xi = (-1)^{\epsilon_\xi \epsilon_\eta + p_\xi p_\eta} \xi \wedge \eta \quad (1.1)$$

inside the exterior algebra. We will often not write the exterior wedges “ $\wedge$ ” explicitly.

2 Antisymplectic Geometry

Consider an antisymplectic manifold (M, E) and let Γ^A denote local coordinates of Grassmann parity $\epsilon_A \equiv \epsilon(\Gamma^A)$ (and exterior form degree $p(\Gamma^A) = 0$). The antisymplectic two-form can locally be written

as

$$E = \frac{1}{2} d\Gamma^A E_{AB} d\Gamma^B = -\frac{1}{2} E_{AB} d\Gamma^B d\Gamma^A , \quad (2.1)$$

where $E_{AB} = E_{AB}(\Gamma)$ is the corresponding matrix representation. Besides carrying gradings $\epsilon(E) = 1$ and $p(E) = 2$, the antisymplectic two-form E has two defining properties. First, E is closed,

$$dE = 0 , \quad (2.2)$$

where the grading conventions for the exterior derivative

$$d = d\Gamma^A \frac{\overrightarrow{\partial}}{\partial \Gamma^A} \quad (2.3)$$

are $\epsilon(d) = 0$ and $p(d) = 1$. Secondly, E is non-degenerate, *i.e.* the antisymplectic matrix E_{AB} has an inverse matrix E^{AB} ,

$$E^{AB} E_{BC} = \delta_C^A = E_{CB} E^{BA} . \quad (2.4)$$

Instead of the compact exterior form notation E , one may equivalently formulate the above conditions with all the indices written out explicitly in terms of the matrices E_{AB} or E^{AB} . In detail, the gradings are

$$\begin{aligned} \epsilon(E_{AB}) &= \epsilon_A + \epsilon_B + 1 = \epsilon(E^{AB}) , \\ p(E_{AB}) &= 0 = p(E^{AB}) , \end{aligned} \quad (2.5)$$

the skew-symmetries are

$$\begin{aligned} E_{BA} &= -(-1)^{\epsilon_A \epsilon_B} E_{AB} , \\ E^{BA} &= -(-1)^{(\epsilon_A + 1)(\epsilon_B + 1)} E^{AB} , \end{aligned} \quad (2.6)$$

while the closeness condition and the equivalent Jacobi identity read

$$\sum_{\text{cycl. } A, B, C} (-1)^{\epsilon_A \epsilon_C} \left(\frac{\overrightarrow{\partial}}{\partial \Gamma^A} E_{BC} \right) = 0 , \quad (2.7)$$

$$\sum_{\text{cycl. } A, B, C} (-1)^{(\epsilon_A + 1)(\epsilon_C + 1)} E^{AD} \left(\frac{\overrightarrow{\partial}}{\partial \Gamma^D} E^{BC} \right) = 0 , \quad (2.8)$$

respectively. The inverse matrix E^{AB} with upper indices gives rise to the antibracket [1]

$$(F, G) = \left(F \frac{\overleftarrow{\partial}}{\partial \Gamma^A} \right) E^{AB} \left(\frac{\overrightarrow{\partial}}{\partial \Gamma^B} G \right) , \quad (2.9)$$

which satisfies a graded skew-symmetry and a graded Jacobi identity as a consequence of eqs. (2.6) and (2.8). There is an antisymplectic analogue of Darboux's Theorem that states that locally there exist Darboux coordinates $\Gamma^A = \{\phi^\alpha; \phi_\alpha^*\}$, such that the only non-vanishing antibrackets between the coordinates are $(\phi^\alpha, \phi_\beta^*) = \delta_\beta^\alpha = -(\phi_\beta^*, \phi^\alpha)$. In Darboux coordinates the antisymplectic two-form is simply $E = d\phi_\alpha^* \wedge d\phi^\alpha$.

3 Odd Laplacian Δ_ρ on Scalars

A scalar function $F = F(\Gamma)$, a density $\rho = \rho(\Gamma)$ and a semidensity $\sigma = \sigma(\Gamma)$ are by definition quantities that transform as

$$F \longrightarrow F' = F , \quad \rho \longrightarrow \rho' = \frac{\rho}{J} , \quad \sigma \longrightarrow \sigma' = \frac{\sigma}{\sqrt{J}} , \quad (3.1)$$

respectively, under general coordinate transformations $\Gamma^A \rightarrow \Gamma'^A$, where $J \equiv \text{sdet} \frac{\partial \Gamma'^A}{\partial \Gamma^B}$ denotes the Jacobian. We shall ignore the global issues of orientation and choice of square root. In principle the above F , ρ and σ could either be bosons or fermions, however normally we shall require the densities ρ to be invertible, and therefore bosons.

Given a choice of density ρ one may define the odd Laplacian [4]

$$\Delta_\rho := \frac{(-1)^{\epsilon_A}}{2\rho} \frac{\overrightarrow{\partial^l}}{\partial \Gamma^A} \rho E^{AB} \frac{\overrightarrow{\partial^l}}{\partial \Gamma^B}, \quad (3.2)$$

that takes scalars to scalars of opposite Grassmann parity. The odd Laplacian (3.2) has a geometric interpretation as a divergence of a Hamiltonian vector field [3, 11]

$$\Delta_\rho \Psi = -\frac{1}{2} \text{div}_\rho(X_\Psi), \quad \epsilon(\Psi) = 1. \quad (3.3)$$

Here $X_\Psi := (\Psi, \cdot)$ denotes a Hamiltonian vector field with a Grassmann-odd Hamiltonian Ψ , and the divergence $\text{div}_\rho X$ of a vector field X , with respect to the measure density ρ , is

$$\text{div}_\rho X := \frac{(-1)^{\epsilon_A}}{\rho} \frac{\overrightarrow{\partial^l}}{\partial \Gamma^A} (\rho X^A), \quad \epsilon(X) = 0. \quad (3.4)$$

The fact that the odd Laplacian (3.3) is non-zero, shows that antisymplectic manifolds do not have an analogue of the Liouville Theorem mentioned in the Introduction. As a consequence of the Jacobi identity eq. (2.8), the square operator $\Delta_\rho^2 = \frac{1}{2}[\Delta_\rho, \Delta_\rho]$ becomes a linear derivation, *i.e.* a first-order differential operator,

$$\Delta_\rho^2(FG) = \Delta_\rho^2(F) G + F \Delta_\rho^2(G). \quad (3.5)$$

Conventionally, one imposes additionally that the Δ_ρ operator is nilpotent $\Delta_\rho^2 = 0$, but this is not necessary for our purposes.

4 Khudaverdian's Δ_E Operator on Semidensities

Khudaverdian showed that one may define a Grassmann-odd, nilpotent, second-order operator Δ_E *without* a choice of density ρ . This Δ_E operator does not take scalars to scalars like the odd Laplacian (3.2), but instead takes semidensities to semidensities of opposite Grassmann parity. Equivalently, the Δ_E operator transforms as

$$\Delta_E \longrightarrow \Delta'_E = \frac{1}{\sqrt{J}} \Delta_E \sqrt{J} \quad (4.1)$$

under general coordinate transformations $\Gamma^A \rightarrow \Gamma'^A$, cf. eq. (3.1). Khudaverdian's construction relies first of all on an atlas of Darboux charts, which is granted by an antisymplectic analogue of Darboux's Theorem, and secondly, on a Lemma by Batalin and Vilkovisky about the possible form of the Jacobians for anticanonical transformations, also known as antisymplectomorphisms.

Lemma 4.1 *“The Batalin-Vilkovisky Lemma” [12, 10, 7, 8, 9, 13]. Consider a finite anticanonical transformation between initial Darboux coordinates $\Gamma_{(i)}^A$ and final Darboux coordinates $\Gamma_{(f)}^A$. Then the Jacobian $J \equiv \text{sdet}(\partial \Gamma_{(f)}^A / \partial \Gamma_{(i)}^B)$ satisfies*

$$\Delta_1^{(i)} \sqrt{J} = 0. \quad (4.2)$$

Here $\Delta_1^{(i)}$ refers to the odd Laplacian (3.2) with $\rho=1$ in the initial Darboux coordinates $\Gamma_{(i)}^A$.

Given Darboux coordinates Γ^A the Δ_E operator is defined on a semidensity σ as [6, 7, 8, 9, 13]

$$(\Delta_E \sigma) := (\Delta_1 \sigma) , \quad (4.3)$$

where Δ_1 is the Δ_ρ operator (3.2) with $\rho=1$. It is important in eq. (4.3) that the formula for the Δ_1 operator (3.2) and the semidensity σ both refer to the same Darboux coordinates Γ^A . The parentheses in eq. (4.3) indicate that the equation should be understood as an equality among semidensities (in the sense of zeroth-order differential operators) rather than an identity among differential operators. One next uses the Batalin-Vilkovisky Lemma to argue that the definition (4.3) does not depend on the choices of Darboux coordinates Γ^A . What this means is, that the right-hand side of the definition (4.3) transforms as a semidensity

$$(\Delta_1^{(f)} \sigma_{(f)}) = \frac{1}{\sqrt{J}} (\Delta_1^{(i)} \sigma_{(i)}) \quad (4.4)$$

under an anticanonical transformation between any two Darboux coordinates $\Gamma_{(i)}^A$ and $\Gamma_{(f)}^A$. Proof:

$$\sqrt{J} (\Delta_1^{(f)} \sigma_{(f)}) = \sqrt{J} (\Delta_J^{(i)} \sigma_{(f)}) = \sqrt{J} (\Delta_J^{(i)} \frac{\sigma_{(i)}}{\sqrt{J}}) = (\Delta_1^{(i)} \sigma_{(i)}) - \frac{1}{\sqrt{J}} (\Delta_1^{(i)} \sqrt{J}) \sigma_{(i)} = (\Delta_1^{(i)} \sigma_{(i)}) . \quad (4.5)$$

The third equality is a non-trivial property of the odd Laplacian (3.2). The Batalin-Vilkovisky Lemma is used in the fourth equality. Strictly speaking, it is enough to consider infinitesimal anticanonical transformations to justify the definition (4.3). The proof of the infinitesimal version of the Batalin-Vilkovisky Lemma goes like this: An infinitesimal anticanonical coordinate transformation $\delta \Gamma^A = X^A$ is necessarily a Hamiltonian vector field $X^A = (\Psi, \Gamma^A) \equiv X_\Psi^A$ with an infinitesimal, Grassmann-odd Hamiltonian Ψ , where $\epsilon(\Psi) = 1$. So

$$\ln J \approx (-1)^{\epsilon_A} (\frac{\vec{\partial}}{\partial \Gamma^A} X^A) = \text{div}_1(X_\Psi) = -2\Delta_1 \Psi , \quad (4.6)$$

and hence

$$\Delta_1 \sqrt{J} \approx -\Delta_1^2 \Psi = 0 , \quad (4.7)$$

due to the nilpotency of the Δ_1 operator in Darboux coordinates. The “ $\approx$ ” sign is used to indicate that equality only holds at the infinitesimal level. (Here we are guilty of mixing active and passive pictures; the active vector field is properly speaking *minus* X .) A simple proof of the Batalin-Vilkovisky Lemma for finite anticanonical transformations can be found in Ref. [13].

On the other hand, once the definition (4.3) is justified, it is obvious that the Δ_E operator super-commutes with itself, because the Γ^A -derivatives have no Γ^A 's to act on in Darboux coordinates. Therefore Δ_E is nilpotent,

$$\Delta_E^2 = \frac{1}{2} [\Delta_E, \Delta_E] = 0 . \quad (4.8)$$

Same sort of reasoning shows that $\Delta_E = \Delta_E^T$ is symmetric.

5 The Δ_E Operator in General Coordinates

We now give a definition of the Δ_E operator that does not rely on Darboux coordinates. We claim that in arbitrary coordinates the Δ_E operator is given as

$$(\Delta_E \sigma) := (\Delta_1 \sigma) + \left(\frac{\nu^{(1)}}{8} - \frac{\nu^{(2)}}{24} \right) \sigma , \quad (5.1)$$

where

$$\nu^{(1)} := (-1)^{\epsilon_A} \left(\frac{\vec{\partial}^l}{\partial \Gamma^B} \frac{\vec{\partial}^l}{\partial \Gamma^A} E^{AB} \right), \quad (5.2)$$

$$\nu^{(2)} := -(-1)^{\epsilon_B} (\Gamma^C, (\Gamma^B, \Gamma^A)) \left(\frac{\vec{\partial}^l}{\partial \Gamma^A} E_{BC} \right) = (-1)^{\epsilon_A \epsilon_C} \left(\frac{\vec{\partial}^l}{\partial \Gamma^A} E^{CD} \right) \left(\frac{\vec{\partial}^l}{\partial \Gamma^D} E^{AB} \right) E_{BC}. \quad (5.3)$$

Eq. (5.1) is the main result of this paper. Notice that in Darboux coordinates, where E^{AB} is constant, *i.e.* independent of the coordinates Γ^A , the last two terms $\nu^{(1)}$ and $\nu^{(2)}$ vanish. Hence the definition (5.1) agrees in this case with Khudaverdian's Δ_E operator (4.3).

It remains to be shown that the right-hand side of eq. (5.1) behaves as a semidensity under general coordinate transforms. Here we will only explicitly consider the case where σ is invertible to simplify the presentation. (The non-invertible case is fundamentally no different.) In the invertible case, we customarily write the semidensity $\sigma = \sqrt{\rho}$ as a square root of a density ρ , and define a Grassmann-odd quantity

$$\nu_\rho := \frac{1}{\sqrt{\rho}} (\Delta_E \sqrt{\rho}) = \nu_\rho^{(0)} + \frac{\nu^{(1)}}{8} - \frac{\nu^{(2)}}{24}, \quad (5.4)$$

by dividing both sides of the definition (5.1) with the semidensity σ . Here we have defined

$$\nu_\rho^{(0)} := \frac{1}{\sqrt{\rho}} (\Delta_1 \sqrt{\rho}). \quad (5.5)$$

Hence, to justify the definition (5.1), one should check that ν_ρ is a scalar under general infinitesimal coordinate transformations. Under an arbitrary infinitesimal coordinate transformation $\delta \Gamma^A = X^A$, one calculates

$$\delta \nu_\rho^{(0)} = -\frac{1}{2} \Delta_1 \text{div}_1 X, \quad (5.6)$$

$$\delta \nu^{(1)} = 4 \Delta_1 \text{div}_1 X + (-1)^{\epsilon_A} \left(\frac{\vec{\partial}^l}{\partial \Gamma^C} E^{AB} \right) \left(\frac{\vec{\partial}^l}{\partial \Gamma^B} \frac{\vec{\partial}^l}{\partial \Gamma^A} X^C \right), \quad (5.7)$$

$$\delta \nu^{(2)} = 3(-1)^{\epsilon_A} \left(\frac{\vec{\partial}^l}{\partial \Gamma^C} E^{AB} \right) \left(\frac{\vec{\partial}^l}{\partial \Gamma^B} \frac{\vec{\partial}^l}{\partial \Gamma^A} X^C \right), \quad (5.8)$$

cf. Appendices A–C. One easily sees that while the three constituents $\nu_\rho^{(0)}$, $\nu^{(1)}$ and $\nu^{(2)}$ separately have non-trivial transformation properties, the linear combination ν_ρ in eq. (5.4) is indeed a scalar.

The new definition (5.1) is clearly symmetric $\Delta_E = \Delta_E^T$ and one may check that the nilpotency (4.8) of the Δ_E operator (5.1) precisely encodes the Jacobi identity (2.8). The odd Laplacian Δ_ρ can be expressed entirely by the Δ_E operator and a choice of density ρ ,

$$(\Delta_\rho F) = (\Delta_1 F) + \frac{1}{\sqrt{\rho}} (\sqrt{\rho}, F) = \frac{1}{\sqrt{\rho}} [\vec{\Delta}_1, F] \sqrt{\rho} = \frac{1}{\sqrt{\rho}} [\vec{\Delta}_E, F] \sqrt{\rho}. \quad (5.9)$$

Since $\nu^{(2)}$ depends on the antisymplectic matrix E_{AB} with lower indices, it is not clear how the formula (5.1) extends to the degenerate anti-Poisson case.

6 Application to Batalin-Vilkovisky Quantization

It is interesting to transcribe the Batalin-Vilkovisky quantization, based on the odd Laplacian Δ_ρ , into a quantization scheme that is based on the Δ_E operator, with the added benefit that no choice of

measure density ρ is needed. Since the Δ_E operator takes semidensities to semidensities, this suggests that the Boltzmann factor $\exp[\frac{i}{\hbar}W_E]$ that appears in the Quantum Master Equation

$$\Delta_E \exp \left[\frac{i}{\hbar} W_E \right] = 0 \quad (6.1)$$

should now be a semidensity, where

$$W_E = S + \sum_{n=1}^{\infty} (i\hbar)^n W_n \quad (6.2)$$

denotes the quantum action. In fact, this was a common interpretation (when restricting to Darboux coordinates) prior to the introduction of a density ρ around 1992, see for instance Ref. [10], p.440-441. If one only considers $\hbar$ -independent coordinate transformations $\Gamma^A \rightarrow \Gamma'^A$ for simplicity, this implies that the one-loop factor e^{-W_1} is a semidensity, while the rest of the quantum action, *i.e.* the classical action S and the higher loop corrections W_n , $n \geq 2$, are scalars as usual. For instance, the nilpotent operator $F \mapsto e^{W_1} \Delta_E (e^{-W_1} F)$ takes scalars F to scalars.

At this stage it might be helpful to compare the above Δ_E approach to the Δ_ρ formalism. To this end, fix a density ρ . Then one can define a bona fide scalar quantum action W_ρ as

$$W_\rho := W_E + (i\hbar) \ln \sqrt{\rho} , \quad (6.3)$$

or equivalently,

$$e^{\frac{i}{\hbar} W_E} = \sqrt{\rho} e^{\frac{i}{\hbar} W_\rho} . \quad (6.4)$$

This scalar action W_ρ satisfies the Modified Quantum Master Equation

$$(\Delta_\rho + \nu_\rho) \exp \left[\frac{i}{\hbar} W_\rho \right] = 0 , \quad (6.5)$$

cf. eq. (5.4), (5.9), (6.1) and (6.4). One may obtain the Quantum Master Equation $\Delta_\rho \exp[\frac{i}{\hbar} W_\rho] = 0$ by additionally imposing the covariant condition $\nu_\rho = 0$, or equivalently $\Delta_E \sqrt{\rho} = 0$. However this step is not necessary.

Returning now to the pure Δ_E approach with no ρ , the finite Δ_E -exact transformations of the form

$$e^{\frac{i}{\hbar} W'_E} = e^{-[\vec{\Delta}_E, \Psi]} e^{\frac{i}{\hbar} W_E} , \quad (6.6)$$

play an important rôle in taking solutions W_E to the Quantum Master Equation (6.1) into new solutions W'_E . It is implicitly understood that all objects in eq. (6.6) refer to the same (but arbitrary) coordinate frame. In general, Ψ is a Grassmann-odd operator that takes semidensities to semidensities. If Ψ is a scalar function (=zeroth-order operator), one derives

$$W'_E = e^{X_\Psi} W_E + (i\hbar) \frac{e^{X_\Psi} - 1}{X_\Psi} \Delta_E \Psi . \quad (6.7)$$

The formula (6.7) is similar to the usual formula in the Δ_ρ formalism [13]. One may check that eq. (6.7) is covariant with respect to general coordinate transformations.

The W - X formulation discussed in Ref. [5] and Ref. [13] carries over with only minor modifications, since the Δ_E operator is symmetric $\Delta_E^T = \Delta_E$. In short, the W - X formulation is a very general field-antifield formulation, based on two Master actions, W_E and X_E , each satisfying a Quantum Master Equation. At the operational level, *symmetric* means that the Δ_E operator, sandwiched between two

semidensities under a (path) integral sign, may be moved from one semidensity to the other, using integration by part. This is completely analogous to the symmetry of the odd Laplacian $\Delta_\rho = \Delta_\rho^T$ itself. The X_E quantum action is a gauge-fixing part,

$$X_E = G_\alpha \lambda^\alpha + (i\hbar)H_E + \mathcal{O}(\lambda^*) , \quad (6.8)$$

which contains the gauge-fixing constraints G_α in involution,

$$(G_\alpha, G_\beta) = G_\gamma U_{\alpha\beta}^\gamma . \quad (6.9)$$

The gauge-fixing functions G_α implement a generalization of the standard Batalin-Vilkovisky gauge-fixing procedure $\phi_\alpha^* = \partial\Psi/\partial\phi^\alpha$. In the simplest cases, the gauge-fixing conditions $G_\alpha = 0$ are enforced by integration over the Lagrange multipliers λ^α . See Ref. [13] for further details on the W - X formulation. The pertinent measure density in the partition function

$$\mathcal{Z} = \int [d\Gamma][d\lambda] e^{\frac{i}{\hbar}(W_E + X_E)} \quad (6.10)$$

is now located inside the one-loop parts of the W_E and the X_E actions. For instance, an on-shell expression for the one-loop factor e^{-H_E} is

$$e^{-H_E} = \sqrt{J \text{sdet}(F^\alpha, G_\beta)} , \quad (6.11)$$

where $J = \text{sdet}(\partial\bar{\Gamma}^A/\partial\Gamma^B)$ denotes the Jacobian of the transformation $\Gamma^A \rightarrow \bar{\Gamma}^A$ and $\bar{\Gamma}^A \equiv \{F^\alpha; G_\alpha\}$. The formula (6.11) differs from the original square root formula [14, 15, 13] by not depending on a ρ density, consistent with the fact that e^{-H_E} is no longer a scalar but a semidensity. We recall here the main point that the one-loop factor e^{-H_E} is independent of the F^α 's and the partition function $\mathcal{Z}$ is independent of the G_α 's in involution, cf. eq. (6.9).

To summarize, the density ρ can altogether be avoided in the field-antifield formalism, at the cost of more complicated transformation rules. We stress that the above transcription has no consequences for the physics involved. For instance, the ambiguity that existed in the density ρ is still present in the choice of W_E and X_E .

ACKNOWLEDGEMENT: The Author thanks I.A. Batalin and P.H. Damgaard for discussions. The Author also thanks the organizers of the workshop ‘‘Gerbes, Groupoids, and Quantum Field Theory, May–July 2006’’ at the Erwin Schrödinger Institute for warm hospitality. This work is supported by the Ministry of Education of the Czech Republic under the project MSM 0021622409.

A Proof of eq. (5.6)

Consider a general (not necessarily infinitesimal) coordinate transformation $\Gamma^A \rightarrow \Gamma'^A$ between an ‘‘unprimed’’ and an ‘‘primed’’ coordinate systems Γ^A and Γ'^A , respectively, cf. eq. (3.1). The primed $\nu_\rho^{(0)}$ quantity (5.5) can be re-expressed with the help of the unprimed coordinates as

$$\nu_{\rho'}^{(0)} := \frac{1}{\sqrt{\rho'}}(\Delta'_1 \sqrt{\rho'}) = \frac{1}{\sqrt{\rho'}}(\Delta_J \sqrt{\rho'}) = \frac{1}{\sqrt{\rho}}(\Delta_1 \sqrt{\rho}) - \frac{1}{\sqrt{J}}(\Delta_1 \sqrt{J}) = \nu_\rho^{(0)} - \nu_J^{(0)} , \quad (A.1)$$

where it is convenient (and natural) to introduce the quantity

$$\nu_J^{(0)} := \frac{1}{\sqrt{J}}(\Delta_1 \sqrt{J}) \quad (A.2)$$

with respect to the unprimed reference system. The third equality in eq. (A.1) uses a non-trivial property of the odd Laplacian (3.2). In the infinitesimal case $\delta\Gamma^A = X^A$, the expression for the Jacobian J reduces to a divergence $\ln J \approx \text{div}_1 X$, and one calculates

$$\delta\nu_\rho^{(0)} = \nu_{\rho'}^{(0)} - \nu_\rho^{(0)} = -\nu_J^{(0)} = -\Delta_1(\ln \sqrt{J}) - \frac{1}{2}(\ln \sqrt{J}, \ln \sqrt{J}) \approx -\frac{1}{2}\Delta_1 \text{div}_1 X, \quad (\text{A.3})$$

which is eq. (5.6).

B Proof of eq. (5.7)

The infinitesimal variation of $\nu^{(1)}$ yields 4 contributions to linear order in the variation $\delta\Gamma^A = X^A$,

$$\delta\nu^{(1)} = -\delta\nu_I^{(1)} - \delta\nu_{II}^{(1)} + \delta\nu_{III}^{(1)} + \delta\nu_{IV}^{(1)}. \quad (\text{B.1})$$

They are

$$\delta\nu_I^{(1)} := (-1)^{\epsilon_A} \left(\frac{\vec{\partial}^l}{\partial\Gamma^B} X^C \right) \left(\frac{\vec{\partial}^l}{\partial\Gamma^C} \frac{\vec{\partial}^l}{\partial\Gamma^A} E^{AB} \right), \quad (\text{B.2})$$

$$\delta\nu_{II}^{(1)} := (-1)^{\epsilon_A} \frac{\vec{\partial}^l}{\partial\Gamma^B} \left(\left(\frac{\vec{\partial}^l}{\partial\Gamma^A} X^C \right) \left(\frac{\vec{\partial}^l}{\partial\Gamma^C} E^{AB} \right) \right) = \delta\nu_I^{(1)} + \delta\nu_V^{(1)}, \quad (\text{B.3})$$

$$\delta\nu_{III}^{(1)} := (-1)^{\epsilon_A} \frac{\vec{\partial}^l}{\partial\Gamma^B} \frac{\vec{\partial}^l}{\partial\Gamma^A} \left(\left(X^A \frac{\overleftarrow{\partial}^r}{\partial\Gamma^C} \right) E^{CB} \right) = \delta\nu_{IV}^{(1)}, \quad (\text{B.4})$$

$$\delta\nu_{IV}^{(1)} := (-1)^{\epsilon_A} \frac{\vec{\partial}^l}{\partial\Gamma^B} \frac{\vec{\partial}^l}{\partial\Gamma^A} \left(E^{AC} \left(\frac{\vec{\partial}^l}{\partial\Gamma^C} X^B \right) \right) = \delta\nu_I^{(1)} + \delta\nu_V^{(1)} + \delta\nu_{VI}^{(1)}, \quad (\text{B.5})$$

$$\delta\nu_V^{(1)} := (-1)^{\epsilon_A} \left(\frac{\vec{\partial}^l}{\partial\Gamma^C} E^{AB} \right) \left(\frac{\vec{\partial}^l}{\partial\Gamma^B} \frac{\vec{\partial}^l}{\partial\Gamma^A} X^C \right), \quad (\text{B.6})$$

$$\delta\nu_{VI}^{(1)} := (-1)^{\epsilon_A} \frac{\vec{\partial}^l}{\partial\Gamma^A} \left(E^{AC} \frac{\vec{\partial}^l}{\partial\Gamma^C} \frac{\vec{\partial}^l}{\partial\Gamma^B} X^B (-1)^{\epsilon_B} \right) = 2\Delta_1 \text{div}_1 X, \quad (\text{B.7})$$

where we have noted various relations among the contributions. Altogether, the infinitesimal variation of $\nu^{(1)}$ becomes

$$\delta\nu^{(1)} = \delta\nu_V^{(1)} + 2\delta\nu_{VI}^{(1)}, \quad (\text{B.8})$$

which is eq. (5.7).

C Proof of eq. (5.8)

The infinitesimal variation of

$$\nu^{(2)} := (-1)^{\epsilon_A \epsilon_C} \left(\frac{\vec{\partial}^l}{\partial\Gamma^D} E^{AB} \right) E_{BC} \left(\frac{\vec{\partial}^l}{\partial\Gamma^A} E^{CD} \right) \quad (\text{C.1})$$

yields 8 contributions to linear order in the variation $\delta\Gamma^A = X^A$, which may be organized as 2×4 terms

$$\delta\nu^{(2)} = 2(-\delta\nu_I^{(2)} - \delta\nu_{II}^{(2)} + \delta\nu_{III}^{(2)} + \delta\nu_{IV}^{(2)}), \quad (\text{C.2})$$

due to a $(A, B) \leftrightarrow (D, C)$ symmetry in eq. (C.1). They are

$$\delta\nu_I^{(2)} := (-1)^{\epsilon_A \epsilon_C} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^D} E^{AB} \right) E_{BF} \left(X^F \frac{\overleftarrow{\partial}^r}{\partial \Gamma^C} \right) \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^A} E^{CD} \right), \quad (\text{C.3})$$

$$\delta\nu_{II}^{(2)} := (-1)^{\epsilon_A \epsilon_C} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^D} E^{AB} \right) E_{BC} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^A} X^F \right) \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^F} E^{CD} \right), \quad (\text{C.4})$$

$$\delta\nu_{III}^{(2)} := (-1)^{\epsilon_A \epsilon_C} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^D} E^{AB} \right) E_{BC} \frac{\overrightarrow{\partial}^l}{\partial \Gamma^A} \left(X^C \frac{\overleftarrow{\partial}^r}{\partial \Gamma^F} E^{FD} \right) = \delta\nu_I^{(2)} + \delta\nu_V^{(2)}, \quad (\text{C.5})$$

$$\delta\nu_{IV}^{(2)} := (-1)^{\epsilon_A \epsilon_C} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^D} E^{AB} \right) E_{BC} \frac{\overrightarrow{\partial}^l}{\partial \Gamma^A} \left(E^{CF} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^F} X^D \right) \right) = \delta\nu_{II}^{(2)} + \delta\nu_{VI}^{(2)}, \quad (\text{C.6})$$

$$\delta\nu_V^{(2)} := (-1)^{\epsilon_A \epsilon_C} E^{FD} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^D} E^{AB} \right) E_{BC} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^A} X^C \frac{\overleftarrow{\partial}^r}{\partial \Gamma^F} \right) = -\delta\nu_V^{(2)} + \delta\nu_{VI}^{(2)}, \quad (\text{C.7})$$

$$\delta\nu_{VI}^{(2)} := (-1)^{\epsilon_A} \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^C} E^{AB} \right) \left(\frac{\overrightarrow{\partial}^l}{\partial \Gamma^B} \frac{\overrightarrow{\partial}^l}{\partial \Gamma^A} X^C \right), \quad (\text{C.8})$$

where we have noted various relations among the contributions. The Jacobi identity (2.8) for E^{AB} is used in the second equality of eq. (C.7). Altogether, the infinitesimal variation of $\nu^{(2)}$ becomes

$$\delta\nu^{(2)} = 3\delta\nu_{VI}^{(2)}, \quad (\text{C.9})$$

which is eq. (5.8).

References

- [1] I.A. Batalin and G.A. Vilkovisky, Phys. Lett. **102B** (1981) 27.
- [2] K. Bering, physics/9711010.
- [3] A. Schwarz, Commun. Math. Phys. **155** (1993) 249.
- [4] I.A. Batalin and I.V. Tyutin, Int. J. Mod. Phys. **A8** (1993) 2333.
- [5] I.A. Batalin, K. Bering and P.H. Damgaard, Phys. Lett. **B389** (1996) 673.
- [6] O.M. Khudaverdian, math.DG/9909117.
- [7] O.M. Khudaverdian and Th. Voronov, Lett. Math. Phys. **62** (2002) 127.
- [8] O.M. Khudaverdian, Contemp. Math. **315** (2002) 199.
- [9] O.M. Khudaverdian, Commun. Math. Phys. **247** (2004) 353.
- [10] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*, Princeton Univ. Press, 1992.
- [11] O.M. Khudaverdian and A.P. Nersessian, Mod. Phys. Lett. **A8** (1993) 2377.
- [12] I.A. Batalin and G.A. Vilkovisky, Nucl. Phys. **B234** (1984) 106.
- [13] I.A. Batalin, K. Bering and P.H. Damgaard, Nucl. Phys. **B739** (2006) 389.
- [14] I.A. Batalin and I.V. Tyutin, Mod. Phys. Lett. **A9** (1994) 1707.

- [15] O.M. Khudaverdian, prepared for the *International Workshop On Geometry And Integrable Models 4-8 October 1994, Dubna, Russia*. In *Geometry and Integrable models*, (P.N. Pyatov and S. Solodukhin, Eds.), World Scientific, 1996.